


# Dijet production at HERA and tests of QCD factorisation



Lidia Goerlich

Institute of Nuclear Physics PAN, Cracow  
on behalf of the H1 and ZEUS collaborations



- Diffraction in ep collisions
- Selection of diffractive events
- Diffractive parton distribution functions
- Exclusive dijet production in diffractive DIS
- Tests of QCD factorisation
- Summary


**MPI @ LHC 2014**

**3-7 November, 2014**

**Kraków, Poland**




# HERA




$E_e = 27.6 \text{ GeV}$

$E_p = 920 - 460 \text{ GeV}$



- HERA – the world’s only ep collider operated in 1992-2007 colliding electrons or positrons with protons
- two colliding beam experiments: H1 and ZEUS
- Nominal proton beam energy :  
 $E_p = 820 / 920 \text{ GeV}$   
 $\sqrt{S} = 300 / 318 \text{ GeV}$ , (HERA- I phase)  
 $E_p = 920 \text{ GeV}$   
 $\sqrt{S} = 318 \text{ GeV}$ , (HERA- II phase)
- Reduced proton beam energy :  
 $E_p = 460 \text{ GeV}$ ,  $\sqrt{S} = 225 \text{ GeV}$ ,  $L_{\text{int}} = 12.4 \text{ pb}^{-1}$   
 $E_p = 575 \text{ GeV}$ ,  $\sqrt{S} = 250 \text{ GeV}$ ,  $L_{\text{int}} = 6.2 \text{ pb}^{-1}$

# Diffraction in ep collisions



Standard DIS variables :

- $Q^2$  |virtuality| of the exchanged boson
- $x$  fraction of proton momentum carried by struck quark in Quark Parton Model
- $y$  inelasticity, fraction of lepton energy transferred in the proton rest frame

$Q^2 \gg 1 \text{ GeV}^2$  deep inelastic scattering (DIS)

$Q^2 \sim 0 \text{ GeV}^2$  photoproduction

**Surprise of HERA** : ~10% of DIS events have no activity in the forward direction ( Large Rapidity Gap)  
 → exchange of a colourless object ( Pomeron, IP)

diffractive variables :

$$x_{IP} = \xi = \frac{Q^2 + M_X^2}{Q^2 + W^2}$$

p-momentum fraction carried by IP


$$\beta = \frac{Q^2}{Q^2 + M_X^2} = \frac{x}{x_{IP}}$$

IP-momentum fraction carried by struck quark

$$t = (p - p')^2$$


squared 4-momentum transfer at proton vertex

# Diffractive dijet production



- Tests of QCD (inspired) models of diffraction
  - Study factorisation properties of diffractive processes
  - Probe partonic structure of diffractive exchange
- Diffractive dijets - direct sensitivity to the gluon component of the Pomeron
- Search for physics beyond DGLAP parton evolution
- Diffraction at HERA – low Bjorken-x phenomenon
- ...

# Factorisation in hard diffraction



**QCD hard scattering collinear factorisation ( proven by Collins 1998) :**

$$d\sigma^{ep \rightarrow eXp}(\beta, Q^2, x_{IP}, t) = \sum f_i^D(\beta, Q^2, x_{IP}, t) \otimes d\sigma^{ei}(\beta, Q^2)$$

$f_i^D$  – diffractive parton density functions (DPDFs), DGLAP evolution in  $Q^2$

$\sigma^{ei}$  – partonic cross sections, same as in inclusive DIS

**Proton vertex factorisation : separate  $(x_{IP}, t)$  from  $(\beta, Q^2)$  dependences ( Ingelman & Schlein, 1985 )**

$$f_i^D(\beta, Q^2, x_{IP}, t) = f_{IP/p}(x_{IP}, t) \cdot F_i^{IP}(\beta, Q^2)$$


**No QCD basis, consistent with experimental data**

**Pomeron flux  
( Regge form )**

**Pomeron  
structure function**



**Diffractive exchange as the Pomeron with a partonic structure**

# Diffractive parton density functions



- Diffractive PDFs obtained through NLO DGLAP QCD fit to data
  - inclusive DDIS cross section → diffractive gluon density weakly constrained at high  $z_{IP}$  → 2 solutions **H1 2006 Fit A and Fit B**
  - combined fit to diffractive inclusive and dijet cross sections → comparable precision of quark and gluon densities for all  $z_{IP}$  (**H1 2007 Jets DPDF, ZEUS DPDF SJ**)

$z_{IP}$  = momentum fraction parton / IP




- Quark Singlet Densities**
- H1 Fit B –  $z\Sigma(z)$
  - H1 Fit Jets –  $z\Sigma(z)$
  - ZEUS SJ –  $z\Sigma(z)\cdot 1.2$
- Gluon Densities**
- H1 Fit B –  $z G(z)$
  - H1 Fit Jets –  $z G(z)$
  - ZEUS SJ –  $z G(z)\cdot 1.2$

**Diffractive scattering is dominated by gluons**

(about 60-70 % of exchanged momentum, extending to large  $z$ )

# Diffractive dijet production


dominant LO QCD diagram  
in diffractive DIS / direct photoproduction



$$x_\gamma = 1$$

$x_\gamma$  – fraction of  $\gamma$  momentum  
in hard subprocess

LO QCD diagram  
in resolved photoproduction  
(  $\gamma$  interacts through its partonic structure )



$$x_\gamma < \sim 0.8$$

Diffractive dijet photoproduction at HERA → sensitive to multi-parton interactions which might occur in resolved photon processes in the presence of photon and Pomeron remnants → breaking of QCD factorisation ?

# Selection of diffractive events


H1 Very Forward Proton Spectrometer

220

H1 Forward Proton Spectrometer

90 80 64 40 24

ZEUS Leading Proton Spectrometer



- **Proton spectrometers:**

- detection of elastically scattered protons → free of proton dissociation background
- low geometrical acceptance → low statistics
- direct measurement of  $t$ ,  $x_{1p}$
- high  $x_{1p}$  accessible

- **Large Rapidity Gap:**

- selection of LRG adjacent to outgoing (untagged) proton
- high acceptance → more statistics
- integration over  $|t| < 1 \text{ GeV}^2$
- background from proton dissociation into low mass resonances  $N^*$


- **The 2 methods have different kinematical coverage, very different systematics**




# Exclusive dijet production in diffractive DIS

New ZEUS analysis of high statistics HERA - II data based on Large Rapidity Gap method

- $e + p \rightarrow e + p + \text{jet} + \text{jet}$
- study of the nature of diffractive exchange
  - investigation of azimuthal angular distribution  
( J. Bartels et al., Phys. Lett. B386 (1996) 389 )




## 2 gluon exchange



Fully perturbative calculations  
based on proton PDF  
( J. Bartels et al. )


## boson-gluon fusion



Resolved Pomeron model  
( Ingelman & Schlein )

# Diffractive dijet production in $\gamma^*$ - IP CMS

**Dijet azimuthal angle  $\phi$**  – angle between the lepton plane and the  $\gamma^*$ - dijet plane in the  $\gamma^*$  - IP rest frame




$$d\sigma/d\phi \propto 1 + A \cos(2\phi)$$


- models predict different shapes for dijet azimuthal angular distribution
- 2g exchange : negative A, maximum of  $d\sigma/d\phi$  at  $\phi = \pi/2$
- boson gluon fusion : positive A, maximum of  $d\sigma/d\phi$  at  $\phi = 0, \pi$

Parton level predictions

**2 gluon exchange**




**boson-gluon fusion**



# Exclusive dijet production in diffractive DIS

New ZEUS analysis of HERA - II data (  $L_{\text{int}} \approx 370 \text{ pb}^{-1}$  ) based on Large Rapidity Gap method


- $Q^2 > 25 \text{ GeV}^2$ ,  $90 < W < 250 \text{ GeV}$   
 $x_{\text{IP}} < 0.01$ ,  $0.5 < \beta < 0.7$   
 $p_{\text{T jet}} > 2 \text{ GeV}$  (  $\gamma^*$ - IP CMS )
- Durham exclusive  $k_t$  jet algorithm :  
final state objects are merged as long as  
 $k_{\text{T}}^2 < y_{\text{cut}} \cdot M_X^2$ ,  
every object must be clustered into a jet
- Jet resolution parameter  $y_{\text{cut}} = 0.15$   
optimizes efficiency vs. purity of dijet  
sample



## SATRAP:

- color dipole model with saturation
- $q\bar{q}$  and  $q\bar{q}g$  in a final state
- good agreement with data
- used in unfolding procedure

## ZEUS



- First measurement of the shape of the azimuthal angular distribution of exclusive dijets in DDIS
- The data favour two gluon exchange model of quark anti-quark production over boson-gluon fusion model

$$d\sigma \propto 1 + A \cos(2\phi), \quad A = -0.18 \pm 0.06(\text{stat.})_{-0.09}^{+0.06}(\text{sys.})$$



# Test of QCD factorisation

Use **HERA DPDFs** and **NLO QCD** calculations to predict diffractive dijet production

Suppression factor  $S^2 = \sigma(\text{data}) / \sigma(\text{theory}_{\text{NLO QCD}})$

- **DIS** : several measurements of H1 and ZEUS → QCD factorisation works within hard diffraction in DIS
- **Factorisation breaking at the Tevatron** ( factor 10 )

Phys. Rev. Lett. 84 (2000) 5043



Similar effect also observed at the LHC  
CMS : Phys. Rev. D87 (2013) 012006


## • Photoproduction :

- H1 : breaking of QCD factorisation ( $S^2 \sim 0.5 - 0.6$ )
- ZEUS: no factorisation breaking ( $S^2 \sim 1$ )

# Diffraction dijet in photoproduction at HERA

Eur. Phys. J. C70 (2010) 15


$E_{\gamma}^{\text{jet1(2)}} > 5(4)$  GeV



$\sigma_{\text{DATA}}/\sigma_{\text{NLO}} \approx 0.6$

Nucl. Phys. B381 (2010) 1

$E_{\gamma}^{\text{jet1(2)}} > 7.5(6.5)$  GeV




$\sigma_{\text{DATA}}/\sigma_{\text{NLO}} \approx 1.0$

- The suppression is expected to be stronger at low scales and low  $x_{\gamma}$
- ... but no evidence for the expected  $x_{\gamma}$  dependence of the suppression factor
- Factorisation breaking observed by H1 but not observed by ZEUS in slightly different phase space

# Diffraction dijet production in DIS


- New H1 analysis of high statistics HERA - II data based on Large Rapidity Gap selection
- $4 < Q^2 < 80 \text{ GeV}^2$ ,  $0.1 < y < 0.7$ ,  $E_{T, \text{jet1(2)}}^* > 5.5(4) \text{ GeV}$ , regularised unfolding procedure



- Data in agreement with the NLO QCD calculations (NLOJET++) using DPDF H1 2006 Fit B
  - confirmation of QCD factorisation
- Data are more precise than theory predictions
  - possible improvement of DPDF fits
  - possibility of determination of the strong coupling constant  $\alpha_s$

# Diffraction dijet production in DIS

## New H1 analysis with proton measured in Very Forward Proton Spectrometer




- Data in agreement with NLO QCD calculations (NLOJET++) using DPDF H1 2006 Fit B → confirmation of QCD factorisation in DDIS
- Photoproduction events selected with same condition except for  $Q^2$

$\gamma p$	DIS
$Q^2 < 2 \text{ GeV}^2$	$4 \text{ GeV}^2 < Q^2 < 80 \text{ GeV}^2$
Common Cuts	
$0.2 < y < 0.7$	
$E_T^{\text{jet1}} > 5.5 \text{ GeV}$	$E_T^{\text{jet2}} > 4.0 \text{ GeV}$
$-1 < \eta^{\text{jet1,2}} < 2.5$	
$0.010 < x_P < 0.024$	
$ t  < 0.6 \text{ GeV}^2$	
$z_P < 0.8$	



# Diffractive dijet in photoproduction


## New H1 analysis with proton measured in Very Forward Proton Spectrometer



- The NLO QCD predictions (Frixione et. al) using DPDF H1 2006 Fit B overestimate the measured cross sections
- No indication of the higher suppression factor at low  $x_\gamma$ , suppression is almost independent on  $x_\gamma$
- Hints for a higher suppression at low  $E_T^{\text{jet1}}$
- Problem of large theoretical uncertainties  $\rightarrow$  use cross section double ratio of data to NLO prediction for photoproduction and DIS

# Proton-tagged diffractive dijets in photoproduction and DIS

The cross section double ratio of data to NLO prediction for photoproduction and DIS



█ █ H1 VFPS Data
 █ █ NLO H1 2006 Fit  $B \times 0.83 \times (1 + \delta_{\text{nadr}})$

Theoretical uncertainties

█ DPDFs uncertainty

█ Overall theoretical uncertainty



- **QCD scale uncertainty:**  
renormalisation and factorisation scales varied simultaneously in photoproduction and DIS by factor of 2 and  $\frac{1}{2}$
- **double ratio of data / NLO**  $\rightarrow$   
theoretical scale uncertainties & most of experimental uncertainties cancel

$$\frac{(\text{DATA}/\text{NLO})_{\gamma P}}{(\text{DATA}/\text{NLO})_{\text{DIS}}} = 0.55 \pm 0.10 (\text{data}) \pm 0.02 (\text{theor.})$$


**Confirmation of QCD factorisation breaking in diffractive dijet photoproduction**  
( suppression not due to proton dissociation )

# Proton-tagged diffractive dijets in photoproduction and DIS

The cross section double ratio of data to NLO prediction for photoproduction and DIS



No dependence of the suppression on  $z_{\text{IP}}$  and  $E_{\text{T}}$  of the leading jet



- **New precise results on diffractive dijets from the H1 and ZEUS Collaborations**
- **The shape of the azimuthal angular distribution of exclusive dijets in diffractive DIS has been measured by ZEUS for the first time**
  - **the data favour 2-gluon exchange model of  $q\bar{q}$  production over boson gluon fusion model**
- **Dijet production with Large Rapidity Gap in diffractive DIS confirms QCD factorisation**
- **New H1 measurements of diffractive dijets in photoproduction and DIS with leading proton :**
  - **suppression factor of 0.55 in photoproduction independent of kinematics is consistent with breaking of QCD factorisation**
  - **the origin of different conclusions of H1 and ZEUS on factorisation breaking in photoproduction not explained**