



Charm and beauty at HERA



Stefan Schmitt, DESY
for the HERA collaborations
H1 and ZEUS




Outline



- The HERA collider
- Charm and beauty production in deep-inelastic scattering
- Data combination
- The new combined HERA charm and beauty data
- Comparisons to NLO QCD
- Charm production in diffractive DIS


- World's only ep collider 1992-2007
- 920 x 27.6 GeV ($\sqrt{s}=320$ GeV)
- Two collider experiments, H1 and ZEUS
- Integrated Luminosity:
 $\sim 2 \times 0.5 \text{ fb}^{-1}$
- e^+p and e^-p data

Two collider experiments:
H1 and ZEUS



Angular coverage with EM+had calorimeters to low angles
Tracking in the central region

- Inclusive processes
 - Neutral current (NC)
 - Charged current (CC)
- Momentum transfer Q^2
- Inelasticity y
- Bjorken- x
- This talk: NC scattering with charm or beauty detected in the hadronic final state X



Measurement at Q^2, x
 probes sum of (anti-) quark PDFs
 $\sigma \sim \sum |M|^2 e_i^2 f_i(Q^2, x)$
 (gluon enters at higher orders)

exchanged 4-momentum:


$$q = e - e' = X - p$$

Kinematic variables

$$Q^2 = -q^2$$

$$y = \frac{p q}{p e}$$

$$x = \frac{Q^2}{s y}$$



Experimental methods:

High p_t lepton
 Reconstructed D, D^* mesons
 Impact parameter, secondary vertex

Measured quantity: reduced cross section σ_{red} with charm or beauty in final state


Reduced cross section: double-differential cross section divided by kinematic factors

NLO calculations: fixed-flavour number scheme (FFNS) where PDF only contains light flavours u, d, s and the gluon. Massive heavy quarks are in the matrix elements


Alternative (not used in this talk): variable-flavour number scheme and massless c, b quarks in the PDF above threshold. PDFs can be converted between schemes.

- Two experiments H1 and ZEUS
 - First combination of HERA charm data published in 2012
 - [Eur.Phys.J.C73 \(2013\) 2311](https://arxiv.org/abs/hep-ex/1207.3434)
 - This talk: new combination of charm and beauty data
 - **H1prelim-17-071, ZEUS-prel-17-01**
 - 13 datasets, using different experimental methods
- (Data combination details: see backup slides)

https://www.desy.de/h1zeus/combined_results/index.php?do=heavy_flavours





Combined data: 12 bins in Q^2 , 52 charm data points and 27 beauty data points, all point-to-point correlations taken into account




+11 more HERA (H1 or ZEUS) analyses

■ H1 VTX ▲ H1 D* HERA-II ◇ ZEUS μ 2005 **H1 and ZEUS preliminary**
□ ZEUS D* 98-00 △ ZEUS D* 96-97 ◇ ZEUS D*
□ ZEUS D* HERA-II ☆ ZEUS VTX ● HERA (prel.)







- Rise to low x : typical for sea and gluon
- Cross section evolves with Q^2
- NLO predictions describe data reasonably well
- Improved precision compared to the 2012 measurement (see backup slides)
- First combination of HERA beauty data




https://www.desy.de/h1zeus/combined_results/index.php?do=heavy_flavours



- Overall satisfactory description of the HERA c and b data by NLO QCD, not much dependent on PDF choice
- No improvement by approximate NNLO
- Slope difference between data and theory as a function of x is visible for charm data at $Q^2 \sim 12 \text{ GeV}^2$



https://www.desy.de/h1zeus/combined_results/index.php?do=heavy_flavours



- Charm and beauty data together with HERA inclusive DIS data are taken as input to a NLO QCD fit (dashed line)

- Simultaneously extract PDFs and c,b masses

$$m_c(m_c) = 1209_{-41}^{+46} (\text{fit})_{-14}^{+62} (\text{model})_{-31}^{+7} (\text{param}) \text{ MeV}$$

$$m_b(m_b) = 4049_{-109}^{+104} (\text{fit})_{-32}^{+90} (\text{model})_{-31}^{+1} (\text{param}) \text{ MeV}$$

- Compatible with previous HERA analyses and with world data

PDG: $m_c(m_c) = 1270 \pm 30 \text{ MeV}$
 and $m_b(m_b) = 4180 \pm 30 \text{ MeV}$


Also see talk by A. Gizhko on running charm mass (Friday)

https://www.desy.de/h1zeus/combined_results/index.php?do=heavy_flavours

- About 10% of the inclusive DIS cross section at HERA are diffractive at low x
- Experimental signature: proton stays intact, no activity in forward detectors, large rapidity gap

t : p vertex 4-mom. transfer squared
 x_{IP} : IP long. mom. fraction
 β or z_{IP} : parton long. mom. fraction


- Theory (Collins): QCD factorisation holds in diffractive DIS \rightarrow concept of diffractive PDFs (DPDFs)



Inclusive diffraction:
extract DPDFs


$$f_i(Q^2, \beta, t, x_{IP})$$

predict \rightarrow




Diffractive charm production: test factorisation theorem in diffraction

DIS phase space
$5 < Q^2 < 100 \text{ GeV}^2$
$0.02 < y < 0.65$
D^* kinematics
$p_{t,D^*} > 1.5 \text{ GeV}$
$-1.5 < \eta_{D^*} < 1.5$
Diffractive phase space
$x_{IP} < 0.03$
$M_Y < 1.6 \text{ GeV}$
$ t < 1 \text{ GeV}^2$



• Electron variables:
 Q^2 and y

• Diffractive variables:
 $\log(x_{IP})$, z_{IP}




- Well described by NLO QCD, large theory scale uncertainties (yellow band)
- DPDF uncertainties (red) similar to data precision
- D^* kinematic distributions also described (\rightarrow backup)


EPJ C77 (2017) 340
[arXiv:1703.09476]

- Investigate diffractively produced fraction of D^* mesons
- Results of many analyses largely agree with each other
- Similar ratios are observed in deep-inelastic scattering and in photoproduction, where one possibly expects to see differences

Note: diffractive QCD factorisation theorem is proven only for DIS [$Q^2 \gg 0$] not for photoproduction [$Q^2=0$]



- New combination of **charm and beauty** double-differential cross section measurements in deep-inelastic scattering at HERA
- Test of QCD with massive quarks (multiple scale problem)
- Fixed flavour-number calculations provide good description
- PDF fit: charm and beauty data constrain quark masses



→ measure running quark masses
from HERA data alone

$$m_c(m_c) = 1209^{+46}_{-41} (\text{fit})^{+62}_{-14} (\text{model})^{+7}_{-31} (\text{param}) \text{ MeV}$$

$$m_b(m_b) = 4049^{+104}_{-109} (\text{fit})^{+90}_{-32} (\text{model})^{+1}_{-31} (\text{param}) \text{ MeV}$$

- New measurement of charm in diffractive DIS at HERA: test of diffractive QCD factorisation and diffractive PDFs
- Data are described by theory within large scale+DPDF uncertainties





Backup


- Two experiments H1 and ZEUS
- First combination of HERA charm data published in 2012
Eur.Phys.J.C73 (2013) 2311
- This talk: new combination of charm and beauty data
H1prelim-17-071, ZEUS-prel-17-01
- 13 datasets, using different experimental methods
(Data combination details: see backup slides)

Data set	Tagging	Q^2 range [GeV ²]	N_c	\mathcal{L} [pb ⁻¹]	\sqrt{s} [GeV]	N_b
1 H1 VTX [8]	VTX	5 – 2000	29	245	318	12
2 H1 D^{*+} HERA-I [9]	D^{*+}	2 – 100	17	47	318	
3 H1 D^{*+} HERA-II (medium Q^2) [10]	D^{*+}	5 – 100	25	348	318	
4 H1 D^{*+} HERA-II (high Q^2) [11]	D^{*+}	100 – 1000	6	351	318	
5 ZEUS D^{*+} 96-97 [12]	D^{*+}	1 – 200	21	37	300	
6 ZEUS D^{*+} 98-00 [13]	D^{*+}	1.5 – 1000	31	82	318	
7 ZEUS D^0 2005 [14]	D^0	5 – 1000	9	134	318	
8 ZEUS μ 2005 [7]	μ	20 – 10000	8	126	318	8
9 ZEUS D^+ HERA-II [2]	D^+	5 – 1000	14	354	318	
10 ZEUS D^{*+} HERA-II [3]	D^{*+}	5 – 1000	31	363	318	
11 ZEUS VTX HERA-II [4]	VTX	5 – 1000	18	354	318	17
12 ZEUS e HERA-II [5]	e	10 – 1000		363	318	9
13 ZEUS μ + jet HERA-I [6]	μ	2 – 3000		114	318	11



Most precise:
secondary
vertex, D^*
meson



- Measurements are extrapolated to a common grid in (Q^2, x) using NLO theory. Correction factors near unity, theory variation considered as systematic uncertainty
- Combination $\chi^2/Ndf=149/187$
- Pull distribution approximately Gaussian




- Many measurements are combined to a single point, large gain in precision
- Correlations of systematic uncertainties between input data points accounted for
- Shown here: charm data before/after combination



Charm: 12 bins in Q^2 , a total of 52 combined data points

Beauty: 12 bins in Q^2 , a total of 27 combined data points

- Compare 2017 combination to 2012 combination of charm data
- Central points are similar
- Improved uncertainties by $\sim 20\%$ at intermediate Q^2





- Calculate chisquare/Ndf between data and theory
- New combination is more precise and has worse chisquare
- The apparent tension is related to shape in x at intermediate Q^2

Dataset	PDF	χ^2	χ^2 with PDF unc.
HERA 2012 c [1] (dof = 52)	HERAPDF20_NLO_FF3A_EIG	59	59
	abm11_3n_nlo	62	62
	ABMP16_3_nnlo	64	63
New combined c (dof = 52)	HERAPDF20_NLO_FF3A_EIG	86	85
	abm11_3n_nlo	92	91
	ABMP16_3_nnlo	101	99
ZEUS VTX b [4] (dof = 17)	HERAPDF20_NLO_FF3A_EIG	14	14
	abm11_3n_nlo	13	13
	ABMP16_3_nnlo	14	14
New combined b (dof = 27)	HERAPDF20_NLO_FF3A_EIG	33	33
	abm11_3n_nlo	34	34
	ABMP16_3_nnlo	39	39

- New analysis by H1
- Large-rapidity gap to select diffractive events
- Electron in backward calorimeter
- D^* reconstructed in $K\pi\pi$ channel
- Cross sections from fit of mass distribution in each analysis bin
- NLO QCD (FFNS) with DPDF from 2006 H1 analysis of inclusive diffraction







D^* in diffractive DIS



About 1100 D^* mesons reconstructed.
Background shape from wrong-charge combinations



DIS phase space
$5 < Q^2 < 100 \text{ GeV}^2$
$0.02 < y < 0.65$
D^* kinematics
$p_{t,D^*} > 1.5 \text{ GeV}$
$-1.5 < \eta_{D^*} < 1.5$
Diffractive phase space
$x_{\mathcal{P}} < 0.03$
$M_Y < 1.6 \text{ GeV}$
$ t < 1 \text{ GeV}^2$

- Analysis of diffractive D^*
- Number of D^* mesons is determined from a fit of the mass in each analysis bin
- The results are well described by the MC model which is used for acceptance corrections



- The cross section is also studied wrt D^* kinematic variables
- The results are described by the NLO calculation

DIS phase space
$5 < Q^2 < 100 \text{ GeV}^2$
$0.02 < y < 0.65$
D^* kinematics
$p_{t,D^*} > 1.5 \text{ GeV}$
$-1.5 < \eta_{D^*} < 1.5$
Diffractive phase space
$x_P < 0.03$
$M_Y < 1.6 \text{ GeV}$
$ t < 1 \text{ GeV}^2$



- D^* production in inclusive DIS has been measured earlier at HERA
- Shown here: ratios of diffractive to inclusive D^* production
- Ratio variations are expected from diffractive phase-space limitations
- Theory: NLO (diffractive) divided by NLO (inclusive) describes data well

DIS phase space
$5 < Q^2 < 100 \text{ GeV}^2$
$0.02 < y < 0.65$
D^* kinematics
$p_{t,D^*} > 1.5 \text{ GeV}$
$-1.5 < \eta_{D^*} < 1.5$
Diffractive phase space
$x_{\mathbb{P}} < 0.03$
$M_Y < 1.6 \text{ GeV}$
$ t < 1 \text{ GeV}^2$

