

# **Analysis of Feynman scaling with the Photon Production in the Very Forward Direction in Deep-Inelastic Scattering at HERA**

## **Abstract**

The production of photons at very small angles with respect to the proton beam direction is studied in deep-inelastic positron-proton scattering at HERA. The data are taken with the H1 detector in the years 2006 and 2007 and correspond to an integrated luminosity of  $126 \text{ pb}^{-1}$ . The analysis covers the range of negative four momentum transfer squared at the positron vertex  $6 < Q^2 < 100 \text{ GeV}^2$  and inelasticity  $0.05 < y < 0.6$ . Cross sections are measured for the most energetic photon with pseudorapidity  $\eta > 7.9$  as a function of Feynman- $x$  variable  $x$  in three intervals of the photon-proton CM energy,  $W$ , in the range  $70 < W < 250 \text{ GeV}$ . The cross sections are normalised to the inclusive deep-inelastic scattering cross section and compared to the predictions of models of deep-inelastic scattering and models of the hadronic interactions of high energy cosmic rays. The fraction of DIS events with forward photons as a function of  $W$  is also measured.





Figure 1: The fraction of DIS events with forward photons in the kinematic region  $6 < Q^2 < 100 \text{ GeV}^2$  and  $0.05 < y < 0.6$  and the pseudorapidity of the photon  $\eta > 7.9$ , as a function of  $W$ . The predictions of MC models are compared to the measurements.






Figure 3: Ratios of normalised cross sections of forward photons production in DIS corresponding to two different  $W$  intervals, shown in Fig.2, as a function of  $x_F$ . The left side figure shows the ratio of the cross section in  $130 < W < 190$  GeV interval to the cross section at  $70 < W < 130$  GeV interval. The right side figure shows the ratio of the cross section in  $190 < W < 250$  GeV interval to the cross section at  $70 < W < 130$  GeV interval. Predictions of LEPTO and CDM MC models are compared to the measurement.




Figure 4: Normalised cross sections of forward photons production in DIS as a function of  $x_F$  in the region  $\eta > 7.9$ ,  $6 < Q^2 < 100 \text{ GeV}^2$  and  $0.05 < y < 0.6$ , and in three  $W$  intervals. The data are compared to predictions of hadronic interactions, QGSJET, EPOS and SIBYLL. The lower row shows the ratios of the MC predictions to the data. The error bars show the total experimental uncertainty, defined as the quadratic sum of the statistical and systematic uncertainties.




Figure 5: Ratios of normalised cross sections of forward photons production in DIS corresponding to two different  $W$  intervals, shown in Fig.4, as a function of  $x_F$ . The left side figure shows the ratio of the cross section in  $130 < W < 190$  GeV interval to the cross section at  $70 < W < 130$  GeV interval. The right side figure shows the ratio of the cross section in  $190 < W < 250$  GeV interval to the cross section at  $70 < W < 130$  GeV interval. Predictions of LEPTO and CDM MC models are compared to the measurement. Predictions of CR models are compared to the measurement.